The Zeldovich Universe Genesis and Growth of the Cosmic Web
Proceedings IAU Symposium No. 308, 2014
Rien van de Weygaert, Sergei Shandarin, Enn Saar & Jaan Einasto, eds.
© 2014 International Astronomical Union
DOI: 00.0000/X000000000000000X

Galaxy and Mass Assembly (GAMA): Selection of the Most Massive Clusters.

Héctor J. Ibarra-Medel[1,2], Maritza Lara-López[3], Omar López-Cruz[1] and the GAMA Team.

[1]National Institute of Astrophysics, Optics and Electronic,
Luis Enrique Erro No 1, Tonantzintla, Puebla, México
email: omarlx@inaoep.mx

[2] email: ibarram@inaoep.mx

[3]Institute of Astronomy, National Autonomous University of México,
Box 70-264, México City, México
email: maritza@astro.unam.mx

arXiv:1410.0748v1 [astro-ph.CO] 3 Oct 2014

Abstract. We have developed a galaxy cluster finding technique based on the Delaunay Tessellation Field Estimator (DTFE) combined with caustic analysis. Our method allows us to recover clusters of galaxies within the mass range of 10^{12} to 10^{16} $\mathcal{M}_{\odot}$. We have found a total of 113 galaxy clusters in the Galaxy and Mass Assembly survey (GAMA). In the corresponding mass range, the density of clusters found in this work is comparable to the density traced by clusters selected by the thermal Sunyaev Zel'dovich Effect; however, we are able to cover a wider mass range. We present the analysis of the two-point correlation function for our cluster sample.

Keywords. methods: data analysis, surveys, galaxies: clusters: general, cosmology: miscellaneous, cosmology: observations

1. Introduction

The GAMA survey (http://www.gama-survey.org/) is a multi-wavelength spectroscopic survey that covers ~ 360 deg^2, which includes $\sim 400,000$ galaxy redshifts down to a magnitude limit of $r_{AB} = 19.8$ (Driver et al. 2011). We chose three stripes within GAMA that cover ~ 144 deg^2 with $\sim 110,000$ galaxy spectra. These three equatorial sky stripes are centred at 9h, 12h and 14.5h (Driver et al. 2011).

We have implemented a new cluster finding technique to find overdensities and estimate cluster masses, simultaneously. We find number galaxy overdensities by using an adaptive method based on the Delaunay Tessellation Field Estimator (DTFE, Schaap & van de Weygaert 2000, Platen 2009), mass estimation is done using caustic analysis (e.g., Serra et al. 2011, Alpaslan et al. 2012). We use this method to detect clusters of galaxies within the mass range of 10^{12} to 10^{16} $\mathcal{M}_{\odot}$, up to $z = 0.3$.

2. Overview and Results

We have found 113 cluster within GAMA. For this sample we have estimated positions, cluster redshifts, velocity dispersions, cluster sizes, and cluster integrated luminosity. Our algorithm has been tested using the GAMA mock catalogs (Robotham et al. 2011). The calculation the cluster luminosities have been generated by using the individual cluster-galaxy luminosity functions (LF) corrected for completeness. We have evaluated the cluster selection function by the application of a simple halo occupation distribution (HOD) model. We want to stress that the density of clusters found by mass selection



Figure 1. The left panel present the comoving density traced by the cluster found in this study (indicated by filled stars), we have compare our results those of Menanteau et al, (2013) generated from a sample of massive clusters selected by the Sunyaev-Zel'dovich effect (open triangles), the continuous dashed line is a simple halo occupation model (HOD). In the right panel we present a compilation of previous results on the characteristic scale for the two-point cluster correlation as a function of cluster separation, the filled big dots represents the results for our sample. We find agreement with previous studies and models

methods (e.g., the Atacama Cosmology Telescope (ACT), Menanteau et al. 2013) is comparable to one found in this work; however, we have covered a larger mass range by more than three orders of magnitude. In addition, we have generated the two-point correlation for clusters of galaxies for our sample. We find broad agreement previous observations and predictions (Estrada et al. 2009). We have generated the mass-to-light ratio (M/L) for the clusters and BCGs in our sample, we find that a single power law $\mathcal{L} \propto \mathcal{M}^{\eta}$ can describe . We found $\eta = 0.6-1$ for clusters and $\eta_{BGC} = 0.1-0.4$ for BCGs. These relations agree with the results of Lin et al.(2004) and Lin & Mohr(2004).

The sample found in this study can be used for further studies in galaxy evolution and its relation with environment. We have shown that optical surveys such as GAMA can be used to select cluster by mass. A sample of cluster selected by our method can be used to traced baryon acoustic oscillations using in a survey in which galaxies are selected in the same fashion as GAMA but covering a larger volume.

References

Alpaslan, M., Robotham, A. S. G., Driver, S., et al. 2012, *MNRAS*, 3012
Bahcall, N. A., Dong, F., Hao, L., et al. 2003, *ApJ*, 599, 814
Driver, S. P., Hill, D. T., et al. 2011, *MNRAS*, 413, 971
Estrada, J., Sefusatti, E., & Frieman, J. A. 2009, *ApJ*, 692, 265
Robotham, A. S. G., Norberg, P., Driver, S. P., et al. 2011, *MNRAS*, 416, 2640
Menanteau, F, et al. 2013, *ApJ*, 765, 67
Platen, E. 2009, Ph.D. Thesis
Schaap, W. E., & van de Weygaert, R. 2000, *A& A*, 363, L29
Serra, A. L., Diaferio, A., Murante, G., & Borgani, S. 2011, *MNRAS*, 412, 800
Lin, Y.-T., Mohr, J. J., & Stanford, S. A. 2004, *ApJ*, 610, 745
Lin, Y.-T., & Mohr, J. J. 2004, *ApJ*, 617, 879
Younger, J. D., Bahcall, N. A., & Bode, P. 2005, *ApJ*, 622, 1